

Gas Dynamics in the Barred Seyfert Galaxy NGC4151 - II. High Resolution HI Study

C.G. Mundell^{1*}, A. Pedlar¹, D.L. Shone¹, A. Robinson²

¹*University of Manchester, Nuffield Radio Astronomy Laboratories, Jodrell Bank, Macclesfield, Cheshire SK11 9DL, UK.*

²*Division of Physical Sciences, University of Hertfordshire, College Lane, Hatfield, Hertfordshire, AL10 9AB, UK.*

**Present address: Department of Astronomy, University of Maryland, College Park, MD 20742, USA.*

Accepted for publication in MNRAS

ABSTRACT

We present sensitive, high angular resolution ($6'' \times 5''$) $\lambda 21$ -cm observations of the neutral hydrogen in the nearby barred Seyfert galaxy, NGC4151. These HI observations, obtained using the VLA in B-configuration, are the highest resolution to date of this galaxy, and reveal hitherto unprecedented detail in the distribution and kinematics of the HI on sub-kiloparsec scales. A complete analysis and discussion of the HI data are presented and the global properties of the galaxy are related to the bar dynamics presented in Paper I.

HI absorption, consistent with previous studies, is detected against the radio continuum nucleus and shows two components – a deep absorption component, centred at $987 \pm 1 \text{ km s}^{-1}$ and width $87 \pm 3 \text{ km s}^{-1}$, and a weaker component, redshifted to $1096 \pm 6 \text{ km s}^{-1}$ with a width of $35 \pm 15 \text{ km s}^{-1}$. An alternative fit is also presented. In addition to the absorption, a high velocity cloud is detected in emission, coincident with the nucleus. This cloud is red-shifted by 260 km s^{-1} from systemic, has an HI mass of $2.3 \times 10^7 M_{\odot}$, and corresponds to outflow on the far side of the nucleus.

Contrary to previous studies, no HI bridge is detected reaching from the shocks directly across the nucleus. Instead, the gas streams from the shocks onto smaller orbits and forms fingers of HI which wind *around* the nucleus, consistent with predictions from general numerical simulations of bars. These fingers correspond closely with dust arcs seen in optical studies and resemble nuclear features seen by others in weak barred galaxies such as M100.

A new rotation curve is presented, extending to within $8''$ of the nucleus and showing a turnover at a radius of $\sim 35''$, which was previously undetected in lower resolution studies. The corresponding resonance curve and the properties of the shocks (Paper I) yield a bar pattern speed of $24.5 \pm 3.7 \text{ km s}^{-1}$ and one Inner Lindblad Resonance (ILR) at a radius of $2.8 \pm 0.6 \text{ kpc}$. Our observations, however, do not rule out the possibility of an Inner ILR.

Key words: galaxies: individual: NGC4151 – galaxies: Seyfert – galaxies: bars – radio lines: atomic

1 INTRODUCTION

Neutral hydrogen is present on a wide range of scales in galaxies, often extending far beyond the optical extent of a galaxy (Broeils & Rhee, 1997 and references therein), and is an important tracer of galactic structure and dynamics. The presence of HI in a galaxy is related to the formation of many observed features; dissipation in the gaseous ISM enables the formation of transient spirals, bars and other large scale structures (Schwarz, 1981; Palouš, 1997). Seyfert galaxies are the closest, most common type of Active Galac-

tic Nuclei (AGN) and are often gas-rich, making them useful laboratories for the study of the AGN phenomenon and its relationship to the gaseous medium in the host galaxy. On the largest scales, HI is sensitive to interactions and tidal features may extend up to 200kpc (e.g. Simkin et al., 1987; Mundell et al., 1995a). On intermediate scales spiral features and galactic bars are well traced in HI and on sub-kpc scales HI may play a role in the fuelling of AGN and in turn be affected by the nuclear activity.

Nuclear activity in galaxies, either as AGN or as starbursts, requires a ready supply of fuel and an efficient mech-

anism for its delivery to the site of activity (Shlosman & Noguchi, 1993). In order to fuel an AGN, the gas must lose sufficient angular momentum to reach the very central accretion radii without triggering star formation, which would inhibit accretion processes (Barnes & Hernquist, 1992). To date this has been simulated quite successfully down to scales of ~ 1 kpc but significant momentum losses on smaller scales have been difficult to achieve.

The rate of accretion or formation and growth of the central black hole itself may be regulated by processes in the host galaxy, such as galactic disk instabilities at different epochs (Shlosman & Noguchi, 1993). Bars, which may be tidally triggered by galaxy interactions (Barnes & Hernquist, 1992) or formed by gravitational disk instabilities, are thought to be an efficient mechanism for transporting galactic material into the central regions of galaxies (Larson, 1994; Wada & Habe, 1995), and for dissipating angular momentum via shocks in the bar (Athanasoula, 1992b). Many nearby barred spiral galaxies possess central mass concentrations with associated non-circular motions and Gerhard (1992) suggests that radial gas inflow is a “common phenomenon”. A subtle balance however, must be maintained such that the fuel is delivered to the AGN without accumulating so large a central mass concentration that the bar itself is destroyed (perhaps 1–5% of the galaxy mass concentrated in a dense core – Hasan & Norman, 1990, Hasan, Pfenniger & Norman, 1993, Friedli, 1994, Sellwood, 1996). The accumulation of a central mass causes the periodic orbits in the bar to become increasingly chaotic, leading to the destruction of the bar and hence the supposed fuelling mechanism.

Statistical studies of the correlation between bars and Seyfert activity are controversial (Arsenault, 1989; Ho, Filippenko & Sargent, 1997; Heckman 1980; Simkin Su and Schwartz, 1980; Moles et al 1995). However, there is increasing evidence from near-infrared studies that as many as 70% of galaxies may possess bars (Mulchaey, Regan and Kundu, 1997), although a significant number of Seyfert galaxies do not (Mulchaey & Regan, 1997). The situation is somewhat complicated with the possibility that those AGN without a bar now, may have had a bar in the past, which has been destroyed by the accumulation of the central mass. (Such Seyferts may lack the gaseous ISM needed to ‘cool’ and maintain the bar). Detailed observations and theoretical studies of bars in AGN complement such statistical studies in an attempt to understand the physical processes at work.

Perhaps one of the most famous and best studied Seyfert galaxies is NGC4151. This galaxy, and its nucleus, has been studied extensively at all wavelengths and was one of the original Seyferts noted in the 1943 sample (Seyfert, 1943) of galaxies with bright star-like nuclei and broad emission-line spectra. NGC4151 has a Seyfert type 1.5 nucleus (Osterbrock and Koski, 1976) in an (R')SAB(rs)ab host galaxy (de Vaucouleurs et al., 1991). Over one thousand papers have been published on NGC4151 and a detailed assessment of the observational status of NGC4151 at all wavelengths has been presented in an excellent review by Schultz (1995).

The most recent classification of NGC4151, that of type (R')SAB(rs)ab (de Vaucouleurs et al., 1991), takes into account the well-defined two-arm outer spiral pattern and the central bulge or ‘fat’ bar. Initially, however, NGC4151 was

believed to be a small spiral galaxy, inclined at 37° with its major axis in PA $\sim 130^\circ$. This description arose because the optical emission originates predominantly from the active nucleus and the central fat bar (a small string of HII regions situated near the each end of the bar looked like small spiral arms). Early HI observations (Davies, 1973), however, revealed a much larger outer spiral structure that suggested a more face-on inclination of $\sim 20^\circ$ and a major axis PA of 26° . These outer spiral arms begin at each end of the bar and wind out to a radius of $\sim 6'$. The spiral arms are particularly gas-rich and, along with the bar, are clearly visible in subsequent HI images by Bosma, Ekers & Lequeux (1977) and Pedlar et al. (1992). The outer arms are faintly discernable in Arp’s (1977) deep optical exposure of the galaxy but it is surprising that such a high concentration of neutral gas produces so little star formation.

NGC4151 is interesting as both an AGN and a barred galaxy and in this paper, we address both the physical properties which relate to the bar and outer spiral structure, and which are of general interest in the study of barred galaxies, and the role which the bar may play in the fuelling of the active nucleus.

We present here a complete analysis and discussion of the highest resolution HI emission-line data of NGC4151 to date. The HI distribution and kinematics of the host galaxy, imaged with the NRAO Very Large Array (VLA)[†] in B configuration, are presented and extend the detailed lower resolution ($17'' \times 22''$) study by Pedlar et al. (1992). Key observational results, which have been described in Paper I (Mundell & Shone, 1998), are related to the global properties of the galaxy, and the implications for fuel transport to the AGN are discussed.

The heliocentric radial velocity of ~ 1000 km s⁻¹ is often used to estimate the distance to NGC4151. However, several authors have noted that by including a correction for the Virgocentric flow, in addition to the galactocentric correction, the distance estimate can increase by almost 50%. Tully (1988) obtained a distance to NGC4151 of 20.3 Mpc ($H_0 = 75$ km s⁻¹ Mpc⁻¹) after applying his Virgocentric correction. More recently, when Aretxaga & Terlevich (1994) took into account the relative velocity of the local group with respect to the Virgo cluster they obtained a corrected recession velocity of 1330 ± 200 km s⁻¹, yielding a distance of 17.7 Mpc ($H_0 = 75$ km s⁻¹ Mpc⁻¹). So for $50 < H_0 < 100$ km s⁻¹ Mpc⁻¹ and $1000 < V < 1523$ km s⁻¹, the distance lies in the range $10 < D < 30.5$ Mpc. The uncertainty in the value of H_0 , however, is as large as the Virgocentric correction (Schultz 1995) so for the purposes of this paper we estimate the distance to NGC4151, using the heliocentric velocity of 998 km s⁻¹ and $H_0 = 75$ km s⁻¹ Mpc⁻¹, to be 13.3 Mpc. One arcsecond corresponds to 65pc in the galaxy at this distance.

2 OBSERVATIONS AND REDUCTION

The $\lambda 21$ -cm radio observations of NGC4151 were obtained in April 1993, using the Very Large Array (VLA) in B con-

[†] NRAO is operated by Associated Universities, Inc., under co-operative agreement with the National Science Foundation.

figuration. Spectral line mode 1A was used, with automatic band-pass calibration and on-line Hanning smoothing performed on the data. Data suffering from ‘shadowing’ (where one antenna’s line of sight is obscured by another) were discarded.

Observations of the point source 1225+368 were interleaved through the source observations throughout the observing run to enable phase calibration of the data. The absolute flux scale was determined from observations of 3C286, assuming a flux density of 14.755 Jy at 1415 MHz (Baars et al. 1977). 3C286 was also used to determine the bandpass corrections in the spectral line data. The observing parameters are given in Table 1.

All data processing was performed using the NRAO AIPS (Astronomical Image Processing System) package. The calibration and initial data editing were carried out using the ‘pseudo-continuum’ channel (channel 0) which is formed on-line by averaging the central 75% of the band (4.7 MHz). The resulting phase and gain solutions were then applied to the B array spectral line data. These data were then Fourier transformed to produce two cubes, one using natural weighting to optimise surface brightness sensitivity, the other uniformly weighted for maximum resolution. The cube dimensions were $512 \times 512 \times 64$, with $2''$ pixels.

Continuum images were formed by averaging 10 channels, free of line emission (channels 10-15 and 55-60), in the natural and uniform cubes, avoiding channels at the ends of the band. The dirty continuum images were subtracted from the corresponding dirty spectral line cubes, to form continuum free cubes.

The continuum images and continuum-free cubes were then deconvolved using the CLEAN algorithm (Högbom, 1974), to remove sidelobe structure. The images produced from the naturally and uniformly weighted data were restored with effective beams of 6×5 arcsec and 4.71×4.71 arcsec half-power full width (FWHM) respectively.

3 RESULTS

3.1 $\lambda 21$ -cm Continuum Emission

The CLEANed $\lambda 21$ -cm continuum image, formed from the line-free channels (see Section 2) in the uniformly weighted cube, is shown in Figure 1. The total flux density from NGC4151 is 336 mJy, consistent with the spectral index of -0.71 found by Hamilton (1995). The rms noise in the image is 0.26 mJy beam $^{-1}$ and the beamsize is $4.71'' \times 4.71''$ (PA 45°). The emission is marginally resolved at this resolution and a two-dimensional Gaussian fit to the component gives dimensions of $5.41'' \times 4.78''$ (PA 79.1°). This slight elongation in PA 79° is consistent with the presence of the ~ 4 -arcsec radio jet in PA 77° , seen at higher resolution and at other frequencies (Pedlar et al., 1993, Mundell et al., 1995b).

3.2 Absorption Against the Radio Continuum Nucleus

Channel maps of the absorption feature, from the continuum subtracted uniform cube, are shown in Figure 1 along with the uniform continuum image in the top left corner. The

Table 1. Observing Parameters for VLA B array observations of NGC4151

GALAXY	NGC4151
Date of Observations	April, 1993
Field Centre (B1950)	$\alpha = 12^h 08^m 01.0^s$ $\delta = 39^\circ 40' 60.0''$
Central Velocity	1000 km s^{-1}
No. of Antennas	27
Time on Source	8.7 hours
Frequency Channels	64
Total Bandwidth	3.125 MHz
Channel Separation	48.828 kHz
Velocity Resolution	12.5 km s^{-1}
Amplitude Calibrator	3C286
Phase Calibrator	1225+368
Spectral Line Mode	1A
Hanning Smoothing	Yes
Primary Beam	$30'$
Cellsize (arcsec)	2
Channel No.s in squash	10-15 & 55-60

absorption appears to be spatially unresolved at the resolution of $4.71'' \times 4.71''$, as the spectra do not show absorption coincident with the eastern extended continuum emission.

The CLEANed continuum image was added back to the CLEANed continuum-free cube, to provide the correct continuum level adjacent to the line in each pixel. The absorption spectra coincident with the nine brightest continuum pixels were selected to form the average absorption spectrum. A linear baseline was fitted and subtracted from each spectrum before weighting each spectrum with its corresponding continuum temperature. These weighted spectra were then averaged together to produce the continuum-weighted average absorption spectrum; the corresponding optical depth spectrum is shown in Figure 2. Table 2 shows the results of fitting two Gaussian profiles to the absorption spectrum. It was possible to obtain convergence, when fitting two Gaussians to the line, in two different ways; (a) a deep narrow Gaussian centred at $(987 \pm 1) \text{ km s}^{-1}$ and a smaller Gaussian redshifted to $(1096 \pm 6) \text{ km s}^{-1}$; (b) a deep narrow component centred at 988 km s^{-1} and a broad shallow component centred at 986 km s^{-1} . Pedlar et al., (1992) also note the presence of two components in their absorption profile, centred at 988 and 1120 km s^{-1} . In both cases, when the Gaussian components were subtracted from the present data, an emission feature, with a peak of $\sim 1.4 \text{ mJy beam}^{-1}$, a FWHM of $\sim 20 \text{ km s}^{-1}$ and centred at $\sim 1050 \text{ km s}^{-1}$, was visible in the residuals.




Figure 1. Maps of the channels containing absorption against the radio continuum nucleus, from the uniformly weighted cube. The $\lambda 21$ -cm continuum image can be seen in the first (top left) panel and the beamsize for all panels is $4.71'' \times 4.71''$ (shown in the lower left corner of the first panel). The continuum contour levels are $(-0.6, 0.6, 1.2, 2.4, 4.8, 9.6, 19.2, 38.4, 76.8)$ mJy beam $^{-1}$. The absorption contour levels are $(-10, -8, -6, -4, -2, 2)$ mJy beam $^{-1}$ and the velocity (in km s $^{-1}$) of each channel map is marked in the top right corner of each map.

Table 2. Gaussian fits to the average absorption spectrum. Two Gaussian profiles were used each time to model the profile and two types of fit, (a) and (b) were equally applicable.

Line Centre (km s $^{-1}$)	FWHM (km s $^{-1}$)	Peak (mJy beam $^{-1}$)
(a) 987 ± 1	87 ± 3	6.6 ± 0.2
(a) 1096 ± 6	35 ± 15	0.9 ± 0.3
(b) 988 ± 12	73 ± 23	5.6 ± 0.6
(b) 986 ± 2	198 ± 6	1.2 ± 0.6

The absorption appears to be spatially unresolved at the resolution of $4.71'' \times 4.71''$, as the spectra do not show absorption against the eastern extended continuum emission. This is consistent with the absorption taking place on scales smaller than the B array beamsize. In fact, Mundell et al. (1995b) studied this strong, unresolved absorption at very high resolution ($0.15''$) with MERLIN and even on these subarcsecond scales, the absorption, which is seen only against the component which is thought to contain the AGN, is marginally resolved.

3.2.1 High Velocity Cloud?

After subtraction of continuum emission, each channel map was examined for signs of residual HI emission. The maps




Figure 2. The average optical depth spectrum over the central beam area.




Figure 3. Spectrum taken over the central pixel of the nuclear region, showing the high velocity emission feature to the left of the central absorption, centred at ~ 1260 km s $^{-1}$.

of these channels show a compact region of emission, coincident with the nucleus, covering a velocity range of 1249 km s $^{-1}$ to 1280 km s $^{-1}$. A spectrum taken over the nuclear region (Figure 3) shows the deep absorption clearly and the emission feature close to the end of the band, centred at ~ 1260 km s $^{-1}$. This feature may be interpreted as a high velocity nuclear cloud, red-shifted by ~ 260 km s $^{-1}$ with respect to systemic, with a column density of 9.6×10^{20} cm $^{-2}$ and hence HI mass of 2.3×10^7 M $_{\odot}$. Since the cloud is coincident with the nucleus but does not show up in absorption and has red-shifted velocities, it would correspond to outflow on the far side of the nucleus.

As this signal is close to the edge of the band and is also spatially coincident with the nucleus, it could be an arti-

fact due to the strong continuum emission from the nucleus. Broader band observations (or indeed observations with the feature placed at the centre of the band) are required to determine whether this feature is a real HI cloud or if it is a baseline problem caused by the strong nuclear continuum source.

3.3 The HI Emission

Although the resolution of the uniformly weighted cube is higher than that of the naturally weighted cube, making it useful for the continuum and absorption study, the sensitivity to HI emission is insufficient to perform a detailed analysis of the HI field; the lowest detectable column density in the uniform cube is $2.8 \times 10^{20} \text{ cm}^{-2}$ compared with $1.32 \times 10^{20} \text{ cm}^{-2}$ in the natural cube. Therefore all subsequent analysis is performed on the naturally weighted cube only.

The channel maps of those channels containing HI emission compare well with those of Pedlar et al. (1992), with emission from both the bar and outer spiral structure visible. The spectra were well approximated by single Gaussian profiles, which enabled moment analysis to be performed on the data. Maps of the channels containing HI emission from the galactic bar are shown in Figure 4. The emission covers the velocity range 907 km s^{-1} to 1083 km s^{-1} . The absorption feature is also visible at the centre of each map.

3.4 HI Moment Maps

3.4.1 Method of Derivation

Line integral profiles, or moments, can be calculated from a spectral line cube. The zeroth moment corresponds to the integrated intensity over velocity, the first moment to the intensity weighted velocity, and the second moment to the intensity weighted velocity dispersion. Several methods exist to derive moment maps of galaxies. The methods range (in effort) from interactively fitting Gaussian profiles to each spectrum at every pixel position across the galaxy, to using a mainly automated routine such as MOMNT in AIPS. Although the former method gives good results it is extremely time consuming, especially for high resolution data, as a typical galaxy may cover a grid of thousands of pixels (e.g. NGC4151 covers $\sim 300 \times 300$ pixels). The latter method is adequate for relatively high signal-to-noise ratio data and to examine, quickly, the main emission features in a galaxy.

The following method (Brinks, private communication) allows more of the signal to be included in the moment analysis without including more noise that will corrupt the analysis. The method combines features of the ‘cut-off’ and ‘window’ methods (Bosma, 1981). The ‘cut-off’ method involves summing (or analysing) all signals above a specified noise limit, in each pixel, through the cube. Using this method alone however can lead to the inclusion of noise spikes in the final map. The ‘window’ method involves examining each channel map and interactively ‘marking’ or drawing around ‘believable’ regions of emission and storing them for later use. The window method has the advantage over the cut-off method, for low signal to noise data, that uncorrelated noise peaks, far away from the real signal and uncorrelated in velocity, are more easily detected and discriminated against (Teuben et al., 1986).

This combination of the two methods is clearly preferable to using MOMNT, which is based on the ‘cut-off’ method only so requires a higher cut-off level to ensure that noise does not corrupt the analysis.

The B-array HI data cube was clipped so that only emission above 2σ was included. Note that the absorption feature at the centre is excluded by this method and hence analysis of the absorption structure was performed separately (see Section 4). The clipped cube was then examined interactively and ‘real’ emission was ‘windowed’ off. Emission was defined to be ‘real’ if it appeared in more than one consecutive channel (i.e. continuity in velocity - at least to a neighbouring channel) and was also present in the C array data of Pedlar et al. (1992).

All emission outside of these windowed regions was blanked out as noise. This blanked cube was then used as the ‘master’ cube or ‘template’. The original cube was masked using the master cube to produce the final blanked cube. XMOM was then used to perform the moment analysis on this cube to produce the zeroth, first and second moments from pixels that had survived the blanking process.

The resulting moment maps are shown in Figure 5. The B array observations reveal structures that are remarkably similar to those found in the previous C array observations, highlighting the power of this type of moment analysis ‘by hand’ for determining the integrated neutral hydrogen distribution for apparently low sensitivity observations. This method was found to be robust when compared with conventional moment analysis (Thean, 1995).

3.4.2 The Moment Maps

The integrated neutral hydrogen image compares well with the features in the optical image of Arp (1977) and the HI image of Pedlar et al. (1992).

We derive an HI mass for the galaxy of $2.3 \times 10^9 M_{\odot}$, of which $5.8 \times 10^8 M_{\odot}$ is in the bar. These lower limits compare well with those of Pedlar et al., (1992) of $3 \times 10^9 M_{\odot}$ and $5 \times 10^8 M_{\odot}$ for the galaxy and bar respectively. Our estimate for the total HI mass is slightly lower, probably due to some missing extended flux resulting from a lack of short spacings, and the bar estimate is higher because the improved resolution means that less of the emission in the bar was lost to the absorption feature in the centre.

There are local minima of the HI column density in the outer regions towards the NE and the SW, between the spiral arms and the bar, and also inside the bar, approximately NW and SE of the central absorption feature. These are real minima and not absorption features, as no nuclear continuum emission is detected in these regions. They were also noted by Pedlar et al., (1992) so are not due to our sensitivity limitations. The density minima along the bar minor axis, between the bar and spiral arms, may be explained by the presence of unstable periodic orbits around the Lagrangian points, L_4 and L_5 (i.e. points at co-rotation, along the bar minor axis) (Contopoulos, 1981; Schwarz, 1984); the lack of stable periodic orbits means that non-periodic orbits have no centres of attraction, so cannot be trapped, thereby causing large stochasticity and depopulation of these regions (Schwarz, 1981; Contopoulos, 1981).

The first moment (Figure 5(b)), representing the inten-




Figure 4. Channel maps of HI emission from the central oval region, from the naturally weighted cube, as in the previous figure. The beam size is $6.07'' \times 5.11''$ and contour levels are $(-10, -5, -0.9, 0.9, 1.8, 2.7)$ mJy beam $^{-1}$, where 1 mJy beam $^{-1}$ corresponds to $N_H = 3.73 \times 10^{20}$ cm $^{-2}$. Each panel is marked with its central velocity (km s $^{-1}$).

sity weighted velocity field, is consistent with that of Pedlar et al. (1992) and is discussed further in Section 3.6.

The second moment corresponds to the intensity weighted dispersion so gives an indication of the variation in line-widths across the galaxy[‡]. The dispersion values in NGC4151 range from ~ 5 to 25 km s $^{-1}$; most of the galaxy

has dispersions of ~ 5 km s $^{-1}$, typical of galactic values (dispersions of 5–8 km s $^{-1}$ —Dickey, Hanson & Helou, 1990), while slightly higher dispersions (10 km s $^{-1}$) are seen in the outer spiral arms in the regions of enhanced emission. The largest dispersions (up to ~ 25 km s $^{-1}$) occur in the bar shocks (see Paper I).

[‡] For a Gaussian line profile the Full Width at Half Maximum is $2\sqrt{2\ln 2} \sigma$ where σ is the second moment.




Figure 5. (a) Zeroth moment map derived using the interactive moment method. The flux cut-off level was $0.7 \text{ mJy beam}^{-1}$ and the beamsize is $6.07'' \times 5.11''$; (b) First moment map derived using the interactive moment method. The flux cut-off level was $0.9 \text{ mJy beam}^{-1}$ and the resolution is the same as in (a). The colour-velocity key is shown at the bottom with the colours labelled in km s^{-1} .

3.5 The Outer Spiral Structure

The two outer spiral arms of NGC4151 are clearly visible in the present observations and can be traced out to $\sim 5.5'$ (21.5 kpc) from the centre, as in the Pedlar et al., (1992) image. The structure shows all of the features visible in the lower resolution data (Pedlar et al., 1992); e.g. the region of enhanced emission that lies $3.5'$ SW of the nucleus in the eastern spiral arm (which is also a location of enhanced optical emission).

3.5.1 An HI Signature of Seyferts?

As noted by Pedlar et al., the spiral arms in NGC4151 are particularly gas-rich but show little evidence of star formation. Emission from the central oval dominates optical images of the galaxy, but in the HI the column densities in the spiral arms are comparable to those in the bar. Although the optical emission in the spiral arms is faint, deep optical images (Arp, 1977) show that the optical emission and HI distribution are closely matched. There appears to be no HI emission beyond the optical extent of the galaxy. This is in contrast to many normal galaxies in which, more usually, the HI extends beyond the optical (Bosma, 1983, Sancisi, 1983, Wevers, 1984, Broeils & Rhee, 1997). Other Seyferts show a similar effect (e.g., NGC1068, Brinks et al. 1994, Brinks & Mundell 1997; NGC4051, Liszt & Dickey; NGC4939, Mundell 1997; NGC5033, Thean et al. 1997), in which the HI and optical emissions are coincident. The only Seyferts in which the HI extends beyond the optical structure (e.g., Mkn348, Simkin et al., 1987; NGC3227, Mundell et al. 1995a; NGC3982, Mundell 1997; NGC5506, Mundell 1997) show clear evidence of tidal disturbance. Alternatively, the close correspondence between the HI and optical extent may be more typical of early-type barred galaxies; studies of NGC1433 (Ryder et al. 1996) and NGC5850 (Higdon et al., 1998) show a close match between the HI and optical structures.

Detailed HI synthesis studies and deep optical imaging of a larger sample of Seyferts and normal galaxies would be required to determine whether this is an HI property specific to Seyferts.

3.5.2 Evidence for Interaction?

Pedlar et al., (1992) note that the spiral structure appears relatively undisturbed and rule out the possibility that NGC4151 has been involved in a (strong) interaction with any of its neighbours. However, the present observations show some evidence of distortion from a symmetrical two-armed spiral pattern and possible evidence of tidal disturbance.

Firstly, the arm segments that lie in the upper half of the image seem to have been displaced upwards. The eastern arm (see Figure 6) is disconnected from the end of the bar compared with the western arm. A more quantitative examination of the spiral pattern can be performed by using the polar diagram in Figure 6. In this image the emission is plotted as a function of azimuthal angle, θ (measured east from the major axis in PA 22°), against distance from the centre of the galaxy. A circle would be represented by a vertical line in the polar plot. The pitch angle of the spiral at any radius




Figure 6. The spiral structure of NGC4151, on the right, is shown in polar co-ordinates on the left. The vertical axis represents azimuthal angle (θ°), measured anticlockwise from the major axis in PA 22° . The linear scale on the horizontal axis represents distance from the centre of the bar. Each segment of the spiral arms can be easily traced in the polar diagram, as indicated by the dotted lines.

r , defined as the tangent to the spiral arm and the tangent to the circle ($r = \text{constant}$) (Binney & Tremaine, 1987), can be easily measured from the polar diagram. Therefore the angle made by the spiral feature and the vertical axis is the pitch angle.

The pitch angle of the western arm remains constant at $\sim 22^\circ$ in the inner regions (i.e. from $100^\circ < \theta < 220^\circ$ or PA of 122° to 242°), close to the southern end of the bar but then begins to decrease, (at $\theta \sim 220^\circ$), to $\sim 3^\circ$. Each arm also is ‘broken’ at about half way along its bend. The western arm is broken but the broken segment appears to follow round continuously, albeit it at a reduced pitch angle. The eastern arm, however, is broken at $\theta \sim 80^\circ$ (PA 102°), with the two segments appearing shifted or ‘dislocated’ from one another. The pitch angle changes from $\sim 27^\circ$ for the segment that starts from the northern end of the bar, to $\sim 16^\circ$ for the second segment that leads to the outer tip of the arm.

Figure 7 shows NGC4151 and some neighbouring galaxies (Bosma et al., 1977) which have systemic velocity values close to that of NGC4151. NGC4156, which is also marked, has been suggested by Arp (1977) to be interacting with NGC4151 despite its rather larger value of systemic velocity of 6750 km s^{-1} . NGC4145, despite its more suitable velocity, seems to be rather far from NGC4151 to be responsible for the observed disturbance. UGC07175 seems another possible candidate along with UGC07188 but interaction simulations are needed to determine whether any of these galaxies could have been responsible for tidally disturbing NGC4151.

3.6 The HI Rotation Curve of NGC4151

To derive the rotation curve of a spiral galaxy to first order, one may fit a curve such as an exponential (linearly increasing to start with then flat thereafter) or Brandt curve (attains a maximum value then declines as Keplerian thereafter) to the velocity field of the galaxy. However, real rota-




Figure 7. The neutral hydrogen distribution of NGC4151 and the positions of three nearby galaxies (marked by crosses), noted by Bosma et al., (1977), with systemic velocities close to that of NGC4151. NGC4156 is also marked, despite its higher systemic velocity. The systemic velocity of each galaxy is also indicated

tion curves deviate from mathematical models due to warps, deviations from circular orbits and irregular mass distributions. Therefore after using a mathematical model of the whole galaxy velocity field to determine the best starting input parameters, it is better to determine the galaxy parameters by fitting curves of constant velocity to a series of concentric annuli. Generally, the radial velocity, V_r , at some azimuthal angle, θ , on an annulus of radius r can be expressed as $V_r = V_s + V_c \sin i \cos \theta$, where V_c is the circular velocity of the annulus and V_s is the systemic velocity of the galaxy. So, for an annulus of radius r , the velocity will vary sinusoidally with angle θ around the annulus, measured from the line of nodes. V_c and i are coupled (Begeman, 1985) but by keeping i fixed, one can solve for V_c and V_s , assuming that V does not change across the annulus.

The rotation curve of NGC4151 was derived from the first moment map using this ‘tilted’ ring model. The width of the annulus ($6''$) was optimised such that it was sufficiently narrow to assume constant velocity across it and wide enough that it contained enough points to enable con-




Figure 8. (a) The HI rotation curve of NGC4151 derived by fitting constant velocity curves to a series of concentric annuli of thickness $6''$. The open circles represent the rotation curve as derived by Pedlar et al. (1992) (b) The change in the position angle of the line of nodes as a function of distance from the centre of the galaxy. Note the clear offset in the line of nodes associated with the bar. Again, the open circles represent the data of Pedlar et al.

vergence of the fit. Assuming that all rings were coplanar (i.e., neglecting any warping), the inclination was held fixed at 21° throughout the fitting, but the PA of the line of nodes, the co-ordinate of the annulus centre and the systemic velocity were allowed to vary. Figure 8(a) shows the resulting rotation curve derived from the B array observations. The first annulus started at a radius of $8''$ from the centre to avoid the central absorption feature and high velocity cloud. Also shown in Figure 8(a) is the rotation curve derived by Pedlar et al. (1992) from their lower resolution data. The two compare well but the rotation curve from the new high resolution data shows a turnover at a radius of $\sim 35''$ which was not evident in the data of Pedlar et al., due to the lower resolution of their observations and the contamination from the central absorption feature.

To first order, the assumption of circular orbits is valid but, as can be seen in Figure 8(b), there are clearly non-circular motions associated with the bar potential as the PA of the line of nodes changes with distance from the centre. The rotation curve will be used later to investigate the

possible existence and positions of resonances in the galaxy (See Section 4).

4 DISCUSSION

A primary aim of detailed studies of barred galaxies is to ascertain such physical properties as the global mass distribution, angular velocity (pattern speed) and degree of central mass concentration of the bar; properties which are not directly observable. However, the general theoretical understanding relating such physical properties of bars to observable phenomena (such as shocks, dust lanes, rings) has become well established in recent years (e.g. Athanassoula, 1992a,b; Lindblad, Lindblad & Athanassoula, 1996).

Analytical and numerical works have predicted a variety of observable phenomena, which, when detected, allow useful constraints to be placed on the physical properties of the bar (e.g. Sanders & Tubbs, 1980; Van Albada & Roberts, 1981; Combes & Gerin, 1985; Athanassoula, 1992a,b; Sellwood & Wilkinson, 1993, and references therein). The gas dynamics of the bar in NGC4151 are discussed in detail in Paper I (Mundell & Shone, 1998); here we relate the bar dynamics to the global properties of the galaxy. In particular, the characteristics of the streaming shocks in NGC4151 provide valuable information about the types and properties of the different families of orbits present in the bar, which in turn are determined by the existence and location of resonances in the galaxy (Teuben et al., 1986).

4.1 Bar pattern speed and ILRs

The new HI rotation curve of NGC4151 (Figure 8(a)) was smoothed (using a box car average of 10 km s^{-1}) and used to produce the resonance curves from derived values of $\Omega = V \text{ (km s}^{-1}\text{)}/R \text{ (kpc)}$ and

$$k = (2\Omega) \sqrt{1 + \frac{R}{2\Omega} \frac{d\Omega}{dR}}.$$

The derived quantities, Ω , $\Omega \pm k/2$, are plotted as a function of radial distance from the centre of the galaxy (defined by the position of the radio continuum peak) in Figure 9.

In order to determine the bar pattern speed and ILR locations, the position of the CR is required. As described in Paper I, simulations of gas flows in weak bars (Athanassoula, 1992b) show that in order to produce such offset shocks as those present in NGC4151, co-rotation is situated just beyond the end of the bar, at a radius of $(1.2 \pm 0.1)a$ (where a is the semi-major axis of the bar). Table 3 shows the corresponding pattern speed of the bar and location of the ILRs for three different positions of CR, derived from the curves in Figure 9. Although the $\Omega - k/2$ shows a small turnover at the inner most radii, the uncertainty in the amplitude here is such that more data points closer to the nucleus are required to confirm whether this is truly a turnover, which will then result in a second ILR.

It should be noted that although the above resonance treatment is a valid first order approximation, the presence of significant non-circular motions and non-linear shock regions in the bar means that it is not possible to derive the radial mass distribution from the rotation curve alone. Full simulations of the gas response to potentials, constrained by




Figure 9. Resonance curves, Ω , $\Omega \pm k/2$ ($\text{km s}^{-1} \text{ kpc}^{-1}$) plotted against radial distance from the centre of the galaxy, R (kpc).

Table 3. Possible bar pattern speeds (Ω_p) and Inner Lindblad Resonance radii for different chosen values of Co-Rotation radius (CR), where a ($99''$) is the bar semi-major axis.

CR	Ω_p ($\text{km s}^{-1} \text{ kpc}^{-1}$)	ILR (kpc)
1.1 a	28.2	2.2
1.2 a	24.5	2.8
1.3 a	20.8	3.4

additional observed properties presented here, are required (Shone & Mundell in preparation).

4.2 Bar-Driven Spiral Pattern in NGC4151?

A simple model (Sanders & Huntley, 1976) in which a potential due to a rigidly rotating bar is introduced to a uniform disk of gas, which has no self-gravity and initially circular orbits, demonstrated that a strong, trailing spiral pattern is produced when the gas reaches steady state. The gas response, to even a weak bar, is spiral – even though the forcing potential is not – and strong, with a density enhancement of as much as a factor of 2.5 in the spiral arms (Binney & Tremaine, 1987). The spiral forms as a consequence of viscosity in the gas and an open spiral structure, such as that in NGC4151, requires only a bar and a dissipative ISM. This bar-driven spiral model would seem promising for NGC4151 with its weak bar potential and greatly enhanced HI spiral arms.

The spiral arms in NGC4151 (as suggested by the de Vaucouleurs classification) are similar to pseudo-rings commonly observed in SB and SAB galaxies (Buta & Crocker, 1991) and are classified as type R'_2 . This classification describes the $\sim 270^\circ$ winding of the spiral arms with respect to the ends of the bar, which results in a doubling of the spiral arms in the two opposing quadrants trailing the bar. The arms clear the opposite side of the bar completely, thus distinguishing it from the more ring-like R'_1 morphology. Such R'_2 pseudo-rings may be the response of galactic disk gas to forcing by a bar (Schwarz, 1981). In this case the OLR is found to lie close to the minor axis of the pseudo-ring, which

for NGC4151 would be at a radius of ~ 16.5 kpc. Constraining the OLR radius to this value and assuming that the spiral and bar pattern speeds are equal, would result in a CR radius of ~ 9 kpc and a pattern speed of ~ 18 km s $^{-1}$.

This CR radius is larger than the observed bar extent in NGC4151, but Schwarz (1981) notes that although bars end at CR, the maximum of the bar may occur at the 4:1 resonance, some way inside CR. However, the bar-driven spiral models also predict that the spiral pattern extends from CR to beyond the OLR; in NGC4151 the spiral pattern, although asymmetric, begins just beyond the ends of the HI bar, which suggests that CR is located here (as assumed in Section 4.1). Therefore, using the CR radius and pattern speed derived in Section 4.1 and assuming the bar and spiral patterns speeds to be the same, the OLR would be situated at a radius of 10-11 kpc, implying that the spiral pattern extends well beyond the OLR.

Alternatively, Sellwood & Wilkinson (1993) point out that there is no direct observational evidence that spiral patterns are driven by their bars, and instead, it is often suggested that the spiral arms have lower pattern speeds than the bar. This may indeed be the case in NGC4151 where, if we assume the spiral pattern to finish at the Outer Lindblad Resonance and CR to lie just beyond the end of the HI bar, we derive a pattern speed for the spiral of < 15 km s $^{-1}$ compared with ~ 25 km s $^{-1}$ for the bar.

4.3 Closer to the nucleus

Theoretical and numerical simulations have shown that galactic bars can drive large inflows of gas towards the nucleus, creating or enhancing a central mass concentration, and possibly holding important implications for the fuelling of AGN and galactic evolution generally (e.g. Simkin, Su & Schwarz, 1980; Shlosman, Frank & Begelman, 1989; Teuben 1996 and references therein). In the lower resolution HI study of Pedlar et al. (1992), the HI in NGC4151 appeared as a continuous neutral hydrogen feature crossing the nucleus, which they suggested was a bridge of HI, originating in the shocks, continuing across the nucleus and resulting in the HI absorption against the radio continuum nucleus. However, further investigation of both the HI emission and absorption at higher angular resolution reveals this interpretation to be unlikely.

The high resolution (0.16'' or 10 pc) MERLIN absorption study (Mundell et al., 1995b) revealed that the HI absorption did not take place against the entire length of the 4''-long radio continuum jet, but instead was localised to the component (C4) which is thought to contain the active nucleus. The absorption was attributed to a 90 M $_{\odot}$ cloud of HI, 1.5pc in extent, which may form part of a nuclear obscuring torus as advocated in Unification Schemes (Antonucci & Miller, 1985; Mundell et al., 1995b).

The high angular resolution HI emission data presented in this paper allows us to resolve individual features in the bar for the first time (e.g. Mundell & Shone, 1998). These data show that the HI does not stream directly from the shocks onto the nucleus but instead loses angular momentum in the shocks and moves onto smaller orbits closer to the nucleus, forming fingers of HI which wind *around* the nucleus. In addition, the HI fingers correspond closely with two arc-like features discovered in the optical colour index

map of Vila-Vilaro, (1995). These two red arcs lie along opposite quadrants of an ellipse, which has a semi-major axis of 18'' and is approximately perpendicular to the bar major axis. Vila-Vilaro et al. attribute the ring to dust extinction of the normal stellar continuum of the galactic bulge, an interpretation supported by the discovery of polarisation due to dichroic absorption around the ellipse by Draper et al. (1992).

Studies of orbits in non-axisymmetric potentials and the resulting gas flows, which predict the formation of shocks, rings and nuclear spiral-like features similar to those seen in NGC4151, help to clarify why the simplistic interpretation of a continuous bridge of HI crossing the nucleus of NGC4151 is not applicable. The shocks in the bar provide direct evidence of large-extent x_2 (perpendicular) orbits in the bar, without which, such curved offset shocks could not form (Athanasoula, 1992b). The dust ellipse may be associated with gas settling onto the x_2 orbits, as in the case of M100 (Knapen et al., 1995). The condition (Athanasoula, 1992b) that at least one ILR is required for the x_2 family to exist (and if two ILRs exist, the x_2 orbits must lie between them), is satisfied in the case of NGC4151, with the dust ellipse lying inside the ILR, whichever of the three positions of CR is used. If a second ILR does exist it must lie closer to the nucleus and therefore would not be detected in the present HI data.

4.3.1 Comparison with M100

Tightly wound spirals or ring-like features have been observed in M100 (Knapen et al., 1995), a grand design two-armed spiral galaxy with a central bar and mild star-formation activity in the nuclear regions. Self-consistent, three dimensional numerical models, with and without star-formation, were used to understand the circum-nuclear structures in M100 (Heller & Shlosman 1994, Knapen et al. 1995). In the model without star-formation, a large-scale pair of leading-edge, trailing shocks formed, consistent with simulations by Athanasoula (1992) and similar to NGC4151 (Mundell & Shone, 1998). In addition, a pair of tightly wound trailing shocks and a smaller pair of leading shocks formed in the central 2.5 kpc of M100. These two shock systems are shown to interact non-linearly, forming two gas circulations, or oval rings, between the two ILRs in the bar (Shlosman, 1996). Evolution of the system, with self-gravity of the gas included, then shows the outer gaseous ring to settle down onto the the lower energy x_2 orbits, inside the OILR, and the inner ring to shrink across the IILR, settling onto smaller x_1 inside the IILR (Shlosman, 1996).

We speculate that a similar situation may be applicable to NGC4151, with the Vila-Vilaro et al. (1995) dust ellipse corresponding to the M100 outer ring, lying inside the OILR and associated with the x_2 orbits in that region. Although an obvious counterpart to the M100-inner ring is not seen in NGC4151, a number of interesting features do exist on smaller scales such as a small blue ring, approximately 2–3'' in extent (Kotilainen & Ward, 1997) and a red compact structure approximately 1 – 2'' long, oriented approximately along the main HI bar (Perez et al., 1989; Terlevich et al., 1991). Suggestions that the red feature could be due to the presence of an obscuring torus (Perez et al., 1989; Terlevich et al., 1991), which collimates the UV radiation responsi-

ble for producing the ENLR (Robinson et al., 1994), would imply that the properties of the active nucleus may be directly related to gaseous conditions in the bar of the host galaxy (Axon & Robinson, 1996). However, high angular resolution optical and infra-red kinematical studies, tracing the rotation curve closer into the nucleus, and determining the kinematics of the nuclear structures, combined with fully self-consistent three-dimensional numerical simulations of the bar would be required to determine the existence of a second ILR and the true nature of the nuclear optical features; contamination from NLR emission however, does cause significant problems in separating rotation from outflow velocities.

5 CONCLUSIONS

- Sensitive, high angular resolution ($6'' \times 5'' - 390$ pc) $\lambda 21$ -cm observations of the neutral hydrogen in the nearby barred Seyfert galaxy, NGC4151, have been presented.

- On the smallest scales, spatially unresolved HI absorption (at a resolution of $4.71''$) against the radio continuum nucleus is detected. Gaussian fits to the two-component absorption spectrum converge with either a deep narrow Gaussian centred at (987 ± 1) km s $^{-1}$ and a smaller Gaussian redshifted to (1096 ± 6) km s $^{-1}$, or a deep narrow component centred at 988 km s $^{-1}$ and a broad shallow component centred at 986 km s $^{-1}$.

- A high velocity cloud is detected in emission, coincident with the nucleus. If this feature is not due to baseline problems, it indicates the presence of a cloud with an HI mass of $2.3 \times 10^7 M_{\odot}$ and red-shifted by 260 km s $^{-1}$ from systemic, corresponding outflow on the far side of the nucleus.

- The HI bridge which was suggested, from lower resolution studies, to originate in the shocks and cross the nucleus directly, is not detected in the present study. Instead, the gas streaming from the shocks forms fingers of HI which wind around the nucleus.

- The HI fingers correspond closely with dust arcs seen in optical studies; these features are consistent with predictions from general numerical simulations of bars, and resemble nuclear features seen by others in weak barred galaxies such as M100.

- The new rotation curve presented extends to within $8''$ of the nucleus and shows a turnover at a radius of $\sim 35''$, which was previously undetected in lower resolution studies.

- The derived resonance curve and the properties of the shocks (Paper I) yield a bar pattern speed of 24.5 km s $^{-1} \pm 3.7$ km s $^{-1}$ and one (ILR) at a radius of 2.8 ± 0.6 kpc. If a second ILR exists closer to the nucleus it would not be detected in the present HI data.

6 ACKNOWLEDGEMENTS

We thank Elias Brinks for useful discussion and the referee, Professor Ron Buta for helpful suggestions. CGM acknowledges a research fellowship from the U.K. Particle Physics and Astronomy Research Council (PPARC).

REFERENCES

- Antonucci, R.R.J. & Miller, J.S., 1985, *ApJ*, 297, 621
 Aretxaga, I. & Terlevich, R.J., 1994, *MNRAS*, 269, 462
 Arp, H., 1977, *ApJ*, 281, 70
 Arsenault, R., 1989, *A&A*, 217, 66
 Athanassoula, E., 1992a, *MNRAS*, 259, 328
 Athanassoula, E., 1992b, *MNRAS*, 259, 345
 Axon, D.J. & Robinson, A., 1996, in ‘Barred Galaxies and Circumnuclear Activity’, eds. Aa. Sandqvist & P.O. Lindblad, Springer, Nobel Symposium 98, p223
 Baars, J.W.M., Genzel, R., Pauliny-Toth, I.I.K., Witzel, A., 1977, *A&A*, 61, 99
 Barnes, J.E. & Hernquist, L., 1992, *ARA&A*, 30, 705
 Begeman, 1985, PhD Thesis, University of Groningen
 Binney, J. & Tremaine, S., 1987, in *Galactic Dynamics*, Princeton University Press, p345
 Bosma, A., Ekers, R.D. & Lequeux, J., 1977, *A&A*, 57,97
 Bosma, A., 1981, *AJ*, 86, 1791
 Bosma, A., 1983, in ‘Internal Kinematics and Dynamics of Galaxies’, ed. E. Athanassoula, Reidel, Dordrecht, p11
 Brinks, E., Skillman, E.D., Terlevich, R.J. & Terlevich, E., 1994, *IAU Symposium* 159, 422
 Brinks, E. & Mundell, C.G., 1997, in ‘Minnesota Lecture Series on Extragalactic HI’, ASP Conference Series, 106, 268
 Broeils, A.H. & Rhee, M.-H., 1997, *A&A*, 324, 877
 Buta, R. & Crocker, D.A., 1991, *AJ*, 102, 1715
 Clements, E.D., 1981, *MNRAS*, 197, 829
 Combes, F. & Gerin, M., 1985, *A&A*, 150, 327
 Contopoulos, G., 1981, *A&A*, 102, 265
 Davies, R.D., 1973, *MNRAS*, 161, 25p
 de Vaucouleurs, G., de Vaucouleurs, A., Corwin, H.G., Buta, R.J., Paturel, G. & Fouque, P., 1991, *Third Reference Catalogue of Bright Galaxies*, Springer-Verlag
 Dickey, J.M., Hanson, M.M. & Helou, G., 1990, *ApJ*, 352, 522
 Draper, P.W., Gledhill, T.M., Scarrot, S.M., Tadhunter, C.N., 1992, *MNRAS*, 257, 309
 Friedli, D., 1994, in ‘Mass-transfer Induced Activity in Galaxies’, ed. I Shlosman, Cambridge University Press, p268
 Gerhard, O.E., 1992, in ‘Unsolved Problems of the Milky Way’, *IAU Symp.* 169, ed. L. Blitz, 79
 Hamilton, N.G., M.Sc. Thesis, Victoria University of Manchester
 Hasan, H. & Norman, C., 1990, *ApJ*, 361, 69
 Hasan, H., Pfenniger, D. & Norman, C., 1993, *ApJ*, 409, 91
 Heckman, T.M., 1980, *A&A*, 87, 152
 Heller, C.H. & Shlosman, I., 1994, 424, 84
 Higdon, J.L., Buta, R.J. & Purcell, G.B., 1998, *AJ*, 115, 80
 Ho, L.C., Filippenko, A.V. & Sargent, W.L.W., 1997, *ApJ*, 487, 591
 Högbom, J.A., 1974, *A&AS*, 15, 47
 Knapen, J.H., Beckman, J.E., Heller, C.H., Shlosman, I. & de Jong, R.S., 1995, *ApJ*, 454, 623
 Kotilainen, J.K. & Ward, M.J., 1997, *A&AS*, 121, 77
 Larson, R.B., 1994, in ‘Mass-Transfer Induced Activity in Galaxies’, ed. I. Shlosman, CUP, p489
 Lindblad, P.A.B., Lindblad, P.O. & Athanassoula, E., 1996, *A&A*, 313, 65L
 Liszt, H.S. & Dickey, J.M., 1995, *AJ*, 110, 998
 Moles et al., 1995, *ApJ*, 438, 604
 Mulchaey, J.S., Regan, M.W. & Kundu, A., 1997, *ApJS*, 110, 299
 Mulchaey, J.S. & Regan, M.W., 1997, *ApJ*, 482, L135
 Mundell, C.G., Pedlar, A., Axon, D.J., Meaburn, J. & Unger, S.W., 1995a, *MNRAS*, 277, 641
 Mundell C.G., Pedlar, A., Baum, S.A., O’Dea, C.P., Gallimore, J.F. & Brinks, E., 1995b, *MNRAS*, 272, 355
 Mundell, C.G., 1997, in ‘Quasar Hosts’, ed. P. Crane, ESO Conference Series, Springer, in press
 Mundell, C.G. & Shone, D.L., 1998, *MNRAS*, in press
 Osterbrock, D.E. & Koski, A.T., 1976, *MNRAS*, 176, 61p

- Palouš, J., 1997, in ‘Minnesota Lecture Series on Extragalactic HI’, ASP Conference Series, 106, 82
- Pedlar, A., Howley, P., Axon, D.J. & Unger, S.W., 1992, MNRAS, 259, 369
- Pedlar, A., Kukula, M.J., Longley, D.P.T., Muxlow, T.W.B., Axon, D.J., Baum, S., O’Dea, C. & Unger, S.W., 1993, MNRAS, 263, 471
- Perez, E., Gonzalez-Delgado, R., Tadhunter, C. & Tsvetanov, Z., 1989, MNRAS, 241, 31p
- Ryder et al. 1996, ApJ, 460, 665
- Robinson, A. et al., 1994, A&A, 291, 351
- Sancisi, R., 1983, in ‘Internal Kinematics and Dynamics of Galaxies’, ed. Athanassoula, E., Reidel, Dordrecht, p55
- Sanders, R.H. & Huntley, J.M., 1976, ApJ, 209, 53
- Sanders, R.H. & Tubbs, A.D., 1980, A&A, 235, 803
- Schultz, H., 1995, *The Seyfert Galaxy NGC4151*, Habilitationsschrift vorgelegt der Fakultät für Physik und Astronomie der Ruhr-Universität Bochum
- Schwarz, M.P., 1981, ApJ, 247, 77
- Schwarz, M.P., 1984, MNRAS, 209, 93
- Sellwood, J.A. & Wilkinson, A., 1993, Rep. Prog. Phys., 56, 173
- Sellwood, J.A., 1996, in ‘Barred Galaxies’, eds. R. Buta, D.A. Crocker & B.G. Elmegreen, IAU Colloquium 157, 259
- Seyfert, C.K., 1943, ApJ, 97, 28
- Shlosman, I., Frank, J. & Begelman, M.C., 1989, Nature, 338, 45
- Shlosman, I. & Noguchi, M., 1993, ApJ, 414, 474
- Shlosman, I., 1996, in ‘Barred Galaxies and Circumnuclear Activity’, eds. Aa. Sandqvist & P.O. Lindblad, Springer, Nobel Symposium 98, p141
- Simkin, S.M., Su, H.J. & Schwarz, M.P., 1980, ApJ, 237, 404
- Simkin, S., van Gorkom, J., Hibbard, J., & Su, H.-J., 1987, Science, 235, 1367
- Terlevich, R., Sánchez Portal, M., Díaz, A.I. & Terlevich, E., 1991, MNRAS, 249, 36
- Teuben, P.J., Sanders, R.H., Atherton, P.D. & Van Albada, G.D., 1986, MNRAS, 221, 1
- Teuben, P.J., 1996, in ‘Barred Galaxies’, eds. R. Buta, D.A. Crocker & B.G. Elmegreen, IAU Colloquium 157, 299
- Thean, A.H.C., 1995, M.Sc. Thesis, Victoria University of Manchester
- Thean, A.H.C., Mundell, C.G., Pedlar, A. & Nicholson, R.A., 1997, MNRAS, 290, 15
- Tully, R.B., 1988, Nearby Galaxies Catalog (Cambridge: Cambridge University Press)
- Van Albada, G.D. & Roberts, W.W., 1981, ApJ, 246, 740
- Vila-Vilaro et al., 1995, A&A, 302, 58
- Wada, K. & Habe, A., 1995, MNRAS, 277, 433
- Wevers, B.H.M.R., 1984, Ph.D. Thesis, University of Groningen